

Corrected  
version  
(Erratum)

# Vegan Diet

## Position of the German Nutrition Society (DGE)

*Margrit Richter, Heiner Boeing, Dorle Grünewald-Funk, Helmut Hesecker, Anja Kroke, Eva Leschik-Bonnet, Helmut Oberritter, Daniela Strohm, Bernhard Watzl for the German Nutrition Society (DGE)<sup>1</sup>*

### Abstract

On the basis of current scientific literature, the German Nutrition Society (DGE) has developed a position on the vegan diet. With a pure plant-based diet, it is difficult or impossible to attain an adequate supply of some nutrients. The most critical nutrient is vitamin B<sub>12</sub>. Other potentially critical nutrients in a vegan diet include protein resp. indispensable amino acids, long-chain n-3 fatty acids, other vitamins (riboflavin, vitamin D) and minerals (calcium, iron, iodine, zinc and selenium). The DGE does not recommend a vegan diet for pregnant women, lactating women, infants, children or adolescents. Persons who nevertheless wish to follow a vegan diet should permanently take a vitamin B<sub>12</sub> supplement, pay attention to an adequate intake of nutrients, especially critical nutrients, and possibly use fortified foods or dietary supplements. They should receive advice from a nutrition counsellor and their supply of critical nutrients should be regularly checked by a physician.

**Keywords:** vegan diet, critical nutrients, vitamin B<sub>12</sub>

is female, young, educated and wealthy, lives in a city and follows a “healthy lifestyle” [1, 2].

### Vegetarian diets

Depending on the form of vegetarianism (♦ Table 1), either selected animal foods (e.g. eggs or milk) are consumed, or animal foods and additives are totally avoided, including foods where animal components are added during the production process. A vegan diet is a very strict form of vegetarian diets. Vegans exclusively consume plant-based foods and reject all kinds of animal foods, sometimes including honey, as well as articles of daily use made of animal body parts – such as fur and leather [2]. The principle reasons for a vegetarian diet are ethical (e.g. rejection of intensive livestock farming), as well as ecological aspects, sustainability and health aspects. According to current knowledge, the risk of nutrition-related diseases is determined by food composition, particularly the balance between animal and plant-based foods, as well as the degree of processing. Epidemiological studies have shown that a high intake of red meat and, particularly, meat products increases the risk of many diseases (e.g. certain cancer sites [5–7]), whereas high levels of dietary fibre-rich cereal products, vegetables and fruit can decrease the risk of many diseases (e.g. cardiovascular diseases [8], and type 2 diabetes mellitus [9]).

## Introduction

### Background

The vegan diet is exclusively restricted to the consumption of plant-based foods. As with other forms of vegetarian diets, it is becoming increasingly popular among the population of the Western world. It is not known precisely how many individuals in Germany adhere to a vegan diet. The data vary between 0.1% and 1% of the population, corresponding to between 81,000 and 810,000 individuals. Excluding animal foods from the diet is usually a conscious and voluntary decision [1, 2]. A vegan diet as part of a “Western” lifestyle differs from a “traditional” plant-based diet, which is mostly practiced in developing countries, where it is often accompanied by restricted food availability and low energy intake, due to low incomes and levels of education [3]. In Western countries, the typical vegetarian

### Citation:

Richter M, Boeing H, Grünewald-Funk D, Hesecker H, Kroke A, Leschik-Bonnet E, Oberritter H, Strohm D, Watzl B for the German Nutrition Society (DGE) (2016) Vegan diet. Position of the German Nutrition Society (DGE). *Ernährungs Umschau* 63(04): 92–102. **Erratum in: 63(05): M262**

This article is available online:  
DOI: 10.4455/eu.2016.021

<sup>1</sup> Complete information on the authors can be found on p. 99.

Vegetarian diets often have a more favourable composition with respect to these foods than the mixed diet which is conventional in Germany considering the supply of nutrients and other beneficial phytochemicals. A meta-analysis of observational studies found that persons adhering to a vegetarian diet exhibited a lower risk of metabolic and cardiovascular diseases than persons not on a vegetarian diet. The risks of ischaemic heart disease and cancer were also lower [10]. However, within the different studies, it is important to consider which reference groups are used and whether other differences in lifestyle were considered in the statistical analysis. There is evidence that some of the investigated groups had very different lifestyles, so that the lower risk of disease is presumably not only due to nutritional differences [11]. In an analysis of two prospective studies, persons adhering to a vegetarian diet (pesco, ovo-lacto vegetarians and vegans) were compared with persons on a high vegetable mixed diet, containing low levels of meat and meat products. There were no differences between the two groups with respect to mortality [12]. These results indicate that we cannot currently assume a health advantage of vegetarians in comparison with persons consuming

a comparable diet with a low level of meat. However, it can be assumed that a plant-based diet (with or without low levels of meat) is associated to a reduced risk of nutrition-related diseases in comparison with the currently conventional German diet.

### Nutrient intake in vegan nutrition

The mixed diet recommended by the DGE contains low levels of meat and meat products, as well as fish. The animal-based foods in this diet contribute to the supply of the nutrients protein resp. essential amino acids, as well as long chain n-3 fatty acids (eicosapentaenoic acid [EPA] and docosahexaenoic acid [DHA]), vitamin D, riboflavin, vitamin B<sub>12</sub> (cobalamin), calcium, iron, iodine, zinc and selenium. As vegetarian diets (♦ Table 1) avoid several or even all of these foods, an adequate supply of these nutrients may be difficult. Adequate supply of these nutrients must be ensured by a well-directed selection of plant-based foods. Alternative sources of nutrients for vegans are summarised in ♦ Table 2.

Only a few studies have investigated the nutritional physiological quality of restrictive diets such as the vegan diet and compared these with other diets [13–15]. Vegeta-

ble-rich diets such as vegetarian diets (♦ Table 1) are associated with good supplies of some vitamins (vitamin C, vitamin E, thiamine and folate [14, 16–23]), minerals (magnesium and potassium [14, 17–19]), dietary fibre and phytochemicals [24, 25]. In comparison to a mixed diet, all vegetarian diets contain, for example, lower levels of saturated fatty acids and cholesterol [14, 24, 25].

The risk of an inadequate supply of nutrients or of nutritional deficiency progressively increases as the selection of foods becomes more restrictive and the diet becomes less varied. This applies in principle to all forms of nutrition. Adequate nutrition can be achieved for vegetarians who eat no meat or meat products, but consume other animal foods, e.g. milk, dairy products and eggs. In a vegan diet, the most critical nutrient is vitamin B<sub>12</sub>. Potentially critical nutrients in a vegan diet moreover include protein resp. indispensable amino acids, as well as long-chain n-3 fatty acids (EPA and DHA), other vitamins (riboflavin, vitamin D<sup>2</sup>) and minerals (calcium, iron, iodine, zinc and selenium)<sup>3</sup> [14, 25, 29].

An overview of the functions and signs of deficiencies for the potentially critical nutrients can be found in the reference values for nutrient intake [30].

	Foods that are eaten	Foods that are avoided
<b>Pesco vegetarians</b>	plant-based foods, fish, eggs, milk and dairy products	Meat and all products derived thereof
<b>Ovo-lacto vegetarians</b>	plant-based foods, eggs, milk and dairy products	meat, fish (including other aquatic animals) and all products derived thereof
<b>Lacto vegetarians</b>	plant-based foods, milk and dairy products	meat, fish (including other aquatic animals), eggs and all products derived thereof
<b>Ovo vegetarians</b>	plant-based foods, eggs	meat, fish (including other aquatic animals), milk and dairy products and all products derived thereof
<b>Vegans</b>	plant-based foods	all animal foods, also honey

Tab. 1: Forms of vegetarianism resp. vegetarian oriented diets (mod. from: [4])

Potentially critical nutrient <sup>a</sup>	Vegetable nutrient source	Comments on the vegan diet <sup>b</sup>
Protein	legumes, nuts, cereals (whole-grain), oil seeds, potatoes specifically combined and consumed over the day (e.g. cereals + legumes, soya products and/or oil seeds).	<ul style="list-style-type: none"> <li>• If a variety of vegetable protein sources, such as cereals, legumes and potatoes, are distributed over the day [83] along with adequate energy intake [84], protein requirements can be covered [85, 86]. It is unclear whether this is also possible for toddlers. According to YOUNG and PELLETT, normal growth is indeed possible in children if they receive an exclusively plant-based diet and if different protein sources are combined [83]. On the other hand, KRAJCOVICOVA-KUDLACKOVA et al. considered that the requirements for indispensable amino acids cannot be exclusively covered by plant protein in phases of high requirements, such as growth, as the protein quality of plant protein is lower than that of animal protein [87].</li> <li>• By well-directed combining different sources of plant protein, the protein quality of daily protein intake can be increased, as can the intake of all indispensable amino acids [88, 89].</li> <li>• Infants and children require relatively more indispensable amino acids in protein than adults do [90]. During growth, it is therefore essential to ensure that intake of protein and of indispensable amino acids as well as energy intake are adequate.</li> </ul>
Long-chain n-3 fatty acids	food fortified with oil from microalgae	<ul style="list-style-type: none"> <li>• With a strict vegan diet, there is hardly any intake of eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA). Moreover, conversion of the n-3 fatty acid <math>\alpha</math>-linolenic acid (ALA; e.g. from flaxseed, walnuts, rape or their oils) to EPA and DHA is limited [91].</li> <li>• Oils from microalgae contain DHA [92–94]. Microalgal oils from the microalgae <i>Ulkenia</i> and <i>Schizochytrium</i> have been approved as Novel Foods [95, 96].</li> </ul>
Vitamin D	Some mushrooms (e.g. common mushrooms, chanterelles), foods fortified with vitamin D	A unique characteristic of vitamin D is that it cannot only be obtained from food, but can also be synthesised by the human body itself if the skin is exposed to sunlight (UVB) [30].
Riboflavin	Oil seeds, nuts, legumes, various types of vegetable (e.g. broccoli, kale) and whole-grain cereals	
Vitamin B <sub>12</sub>	Foods fortified with vitamin B <sub>12</sub>	<ul style="list-style-type: none"> <li>• If fortified foods are strictly excluded in a vegan diet, hardly any vitamin B<sub>12</sub> is consumed.</li> <li>• As a result of bacterial fermentation, plant-based foods, such as sauerkraut, may contain traces of vitamin B<sub>12</sub>. However, it is unclear whether this form of vitamin B<sub>12</sub> can be utilised in the human body. Moreover, the levels are so low that adequate intake is not possible [2].</li> <li>• Shiitake mushrooms contain vitamin B<sub>12</sub>, although the quantities may vary greatly [97]. Sea algae such as nori may also contain vitamin B<sub>12</sub>. Attention must then be paid to the declared moderate content of iodine. However, these foods are unsuited to be the sole source of vitamin B<sub>12</sub>, as the bioavailability is unclear resp. the vitamin B<sub>12</sub> is in an inactive form [97–102].</li> <li>• Spirulina and other products with cyanobacteria marketed as natural sources of vitamin B<sub>12</sub> for persons adhering to a vegan diet contain no cobalamin in a form that is active in the human body and are therefore unsuited for fulfilling requirements [102].</li> <li>• Vegans cannot ensure their supply of vitamin B<sub>12</sub> with conventional foods, including fermented foods. An adequate supply of vitamin B<sub>12</sub> can only be ensured by taking a vitamin B<sub>12</sub> supplement [32, 33].</li> </ul>
Calcium	Vegetables (e.g. broccoli, kale, rocket), nuts (e.g. hazelnuts and Brazil nuts), legumes, soya meat replacement products (“textured soya protein”), tofu, mineral water (calcium-rich, > 150 mg/L), foods fortified with calcium	
Iron	legumes, oil seeds, nuts, whole-grain cereals and various types of vegetable (e.g. spinach, black salsify)	<ul style="list-style-type: none"> <li>• Simultaneous consumption of foods rich in vitamin C or other organic acids improves iron bioavailability [2].</li> <li>• Substances such as phytates and polyphenols (e.g. in tea and coffee) may reduce iron absorption [2, 103]. Black tea and coffee should not be drunk directly before, during or after meals of high iron content.</li> <li>• As generally recommended, it is particularly important that pregnant women and lactating women should take an iron supplement if they are known to suffer from iron deficiency.</li> </ul>
Iodine	<ul style="list-style-type: none"> <li>• Iodinated and fluoridated table salt as well as foods prepared thereof (as generally recommended)</li> <li>• Sea salt fortified with sea algae and of defined iodine content, or occasional consumption of sea algae of moderate iodine content, e.g. nori</li> </ul>	<ul style="list-style-type: none"> <li>• The Federal Institute for Risk Assessment considers that dried algal products of iodine content &gt; 20 mg/kg are harmful to health and advises against their consumption [104].</li> <li>• Goitrogenic substances in plant-based foods, such as cabbage plants, soya beans and sweet potatoes, may decrease the bioavailability of iodine, which may be important if iodine intake is very low [2].</li> <li>• As generally recommended, it is particularly important that pregnant women and lactating women should take iodine supplements, after consulting a physician and considering their individual iodine intake.</li> </ul>
Zinc	Whole-grain, legumes, oil seeds, nuts	Preparation procedures such as sour dough fermentation and germination improve bioavailability [2].
Selenium	Cabbage (e.g. broccoli, white cabbage), bulb vegetables (e.g. garlic, onions), mushrooms, asparagus and legumes, Brazil nuts	The content in plant-based foods is highly dependent on the area of culture, as it depends on the selenium content in the soil [2].

◀ **Tab. 2: Potential critical nutrients in a vegan diet and vegetable nutrient sources**

<sup>a</sup>For reference values for the intake of these nutrients, see [30] and URL: [www.dge.de/wissenschaft/referenzwerte/](http://www.dge.de/wissenschaft/referenzwerte/)

<sup>b</sup>Aside from the comments here on taking dietary supplements, vegans, like the general population, should comply with the recommendations on vitamin K administration to neonates (oral administration of 3 x 2 mg vitamin K) and on vitamin D administration to infants (daily 10 µg vitamin D) and for the intake of folic acid supplements by women intending to get pregnant (daily 400 µg synthetic folic acid); folic acid intake should start not later than 4 weeks before the start of the pregnancy and continue until the end of the first trimester of pregnancy [30].

## Vitamin B<sub>12</sub>

Vitamin B<sub>12</sub> is solely produced by microorganisms. The form that is available to the human body occurs almost exclusively in animal foods (♦ Table 2) [2]. Thus a vegan diet increases the risk of vitamin B<sub>12</sub> deficiency (unless a vitamin B<sub>12</sub> supplement is used) [31]. Several studies on vegans who took no vitamin B<sub>12</sub> supplements have found that the prevalence of low vitamin B<sub>12</sub> supply resp. vitamin B<sub>12</sub> deficiency was up to 86%, depending on the parameters examined [32, 33].

Vitamin B<sub>12</sub> participates in the regulation of homocysteine concentrations and methionine metabolism, as well as haematopoiesis and the metabolism of fatty acids and amino acids. Low serum concentrations of vitamin B<sub>12</sub> are associated with high plasma concentrations of homocysteine. High plasma homocysteine concentrations have been considered to be a risk factor for cardiovascular diseases [34–36]. Protracted low vitamin B<sub>12</sub> intake can also impair cell function and DNA synthesis and may lead to megaloblastic anaemia or neurological disorders [37].

As the liver store of vitamin B<sub>12</sub> is relatively large and the reutilisation rate through enterohepatic circulation is high, clinical symptoms of deficiency only become evident after some years of vitamin B<sub>12</sub>-free nutrition. Thus, vegans should regularly have their vitamin B<sub>12</sub> supply checked. Neonates depend on the mother receiving adequate vitamin B<sub>12</sub> intake during pregnancy and lactation. There is preliminary evidence that vitamin B<sub>12</sub> deficiency before con-

ception is an independent risk factor for complications during pregnancy (miscarriage, pre-eclampsia) and for negative effects on the neonate (low birth weight, neural tube defect) [38–41]. If the mother is on a strict vegan diet without dietary supplements and the neonate is exclusively breast-fed, vitamin B<sub>12</sub> deficiency and its consequences (neurological disorders, megaloblastic anaemia) could already be observed in the first months of life [42–54].

Vitamin B<sub>12</sub> supply can be assessed with several biomarkers. These include blood concentrations of vitamin B<sub>12</sub>, holotranscobalamin (HTC), serum methylmalonic acid (MMA) and plasma homocysteine. None of these biomarkers is suitable as the sole parameter to evaluate vitamin B<sub>12</sub> supply [55].

### Population groups with special requirements for nutrient supply

In persons on a vegan diet, there are difficulties with the supply of some nutrients (♦ Table 2), and this may have unfavourable consequences for health, particularly for people in sensitive phases of life (e.g. growth). Therefore special attention is required for pregnant and lactating women on a vegan diet, as well as children from infants through the growth phase up to adolescence who adhere to or who are given a vegan diet. These special population groups are at greater risk of nutritional deficiencies [30]. There are currently no conclusive studies with persons on a vegan diet that might allow a comprehensive evaluation of the supply situation

for these groups at risk. Case reports show that the supply of vitamin B<sub>12</sub> and iodine of infants whose mothers were on a vegan diet was not adequately guaranteed and that they developed neurological disorders and megaloblastic anaemia or goitre [46, 47, 52, 56].

Nutrient supply in a vegan diet in pregnancy was investigated in a systematic review. The studies were heterogenous, with respect to both their methodology and their results. Some of the studies were rated as being of poor quality. Bearing in mind these limitations, the authors concluded that a freely chosen vegetarian or vegan diet without financial restrictions is safe during pregnancy, if nutrient supply is observed and nutrient deficits are compensated for [3].

For population groups with special requirements for nutrient supply, a well-directed selection of foods and modes of preparation is necessary, i.e. high quality, nutrient dense foods, preparation and administration appropriate to age. Particularly in phases of growth or of increased requirements for nutrients, the intake and supply of nutrients should be regularly checked by a nutrition counselor or physician. If intake is inadequate, this can be optimised by

<sup>2</sup> Aside from vitamin D intake in food, vitamin D can also be produced in the human body by UVB light exposure (sunlight). The synthesis of vitamin D in skin depends on various factors, e.g. the dose and wavelength of UVB exposure and the area of the exposed skin, the skin type and the age [26, 27].

<sup>3</sup> Among these vitamins and minerals, vitamin D and iodine should be regarded as critical nutrients in the general population, iron and calcium only in specific population groups [28].

increasing the intake of individual nutrients by means of the – precisely calculated – consumption of fortified foods or by taking dietary supplements [2, 3]. According to the current state of knowledge, it is not possible to ensure an adequate vitamin B<sub>12</sub> supply with a vegan diet without taking dietary supplements.

### Food selection in a vegan diet

The foods consumed in a vegan diet are not necessarily nutritionally favourable or health-promoting. The foods consumed may include vegetables, legumes, fruit, nuts, seeds, valuable plant oils or whole-grain products, all of which have been shown to have favourable effects [8, 9, 57]. If however high levels of sugar, fat or table salt are added to vegan dishes, then they are nutritionally unfavourable [9, 57, 58]. Food selection may depend on the individual motive for a vegan diet (e.g. animal welfare or health reasons) [15, 59].

Due to many changes and an enlargement in the available (processed) foods and possible food supplies for vegans in the recent years, the supply situation for persons on a vegan diet has probably improved. About 90% of the 852 vegans surveyed in an online survey (July/August 2013) reported that it had become much simpler in recent years to follow a vegan diet [1]. Due to the increased demand, an abundant range of vegan convenience and replacement products are now available, even including imitations of meat products and cheese. These are intended to respond to the customers' wish for a wide range of choices and accustomed dishes. The manufacturing conditions have long achieved the level for conventional foods [60]. Some of these are highly processed products containing many additives, and their nutritional value has sometimes

been criticised [61–63]. However, these foods can support nutrient supply to vegans, in so far as they are fortified with nutrients.

◆ Table 2 shows the options for vegan nutrition that ensure the intake of potentially critical nutrients. The sources for potentially critical nutrients are natural foods rich in these nutrients, fortified foods and dietary supplements. Vegans should specifically incorporate these in their menus. In this way and with specific food selection and good planning, it is possible to create a vegan diet in which no nutrient deficiency develops.

One sensitive population group are infants who are not – or only partially – breastfed. With industrially produced infant formulas based on cows' milk, healthy non-breastfed infants after a mature birth are provided with adequate nutrients in the first months, independently of the mother's diet. Apart from infant formulas based on cows' milk, there are also infant formulas based on soya. A current review [64] concludes that modern infant formulas based on soya are safe and that their effects on growth, bone health, and on reproductive, endocrine and neurological function and the immune system do not differ from those of other infant formulas or of breast milk. On the other hand, the Federal Institute for Risk Assessment (BfR) [65], the Young Family Network [66] and the Nutrition Committee of the German Paediatric [67, 68] consider that infant formulas based on soya cannot replace products based on cows' milk and are not intended for the nutrition of healthy infants. Because of their content of phyto-oestrogens and the higher level of aluminium in comparison to infant formulas based on cows' milk, infants who have not been breastfed (or only partially) should only be given soya products regularly in exceptional and justified cases (e.g. galactosaemia) and on medical recommendation [65–68].

### Food-related nutrition recommendations

Food-related nutrition recommendations for vegan nutrition have been published by various scientific societies and experts. The Giessen vegetarian food pyramid contains all food groups for ovo-lacto vegetarians and also for vegans, with the comment that vegans should use fortified foods or dietary supplements to cover their requirements [2]. In order to support the planning of a vegetarian diet, the Dietary Guidelines for Australia contain specific notes for ovo-lacto vegetarians and vegans on individual food groups [69]. As part of the 2015 Dietary Guidelines for Americans, the United States Department of Agriculture (USDA) developed nutrition recommendations for a Healthy Vegetarian Pattern [70] and provides suggestions for consumers who wish to adhere to a vegetarian diet [71].

In the Giessen vegetarian food pyramid, it is recommended that vegetarians and vegans should replace meat, fish, milk and eggs with legumes, such as peas, beans, chickpeas and lentils, or with soya products and other protein sources. A comparison between a wholesome diet in accordance with the recommendations of the DGE and the recommendations for a vegan diet in the Giessen vegetarian food pyramid shows that both have the same basis and the corresponding food-related recommendations are very similar (◆ Table 3). However, the following specific points must be considered for vegan nutrition:

**Vegetables, including legumes and fruit** provide the basis for health-promoting nutrition, because of their high nutrient density, their high content of dietary fibre and phytochemicals and their potential to prevent various nutrition-related diseases. “5 a day”, i.e. 5 portions of vegetables and fruit a day, are recommended. For vegetarian (including vegan) nutrition, the Giessen vegeta-

Food	Wohlesome diet (mixed diet) in accordance with the DGE (reference values for adults) [80, 81]	Vegan nutrition in accordance with the Giessen vegetarian food pyramid [2]
<b>Group 1:</b> Cereals, cereal products, potatoes	daily <ul style="list-style-type: none"> <li>4–6 slices (200–300 g) of bread <b>or</b> 3–5 slices (150–250 g) of bread and 50–60 g cereal flakes and</li> <li>1 portion (200–250 g) potatoes (cooked) <b>or</b> 1 portion (200–250 g) noodles (cooked) <b>or</b> 1 portion (150–180 g) rice (cooked)</li> </ul> Preferably whole-grain products	About 2–3 meals a day, e.g. whole-grain bread, potatoes, rice
<b>Group 2:</b> Vegetables (including legumes) and salad	daily <ul style="list-style-type: none"> <li>at least 3 portions (400 g) of vegetables: 300 g cooked vegetables and 100 g raw fruit and vegetables/salad <b>or</b> 200 g cooked vegetables and 200 g raw fruit and vegetables/salad</li> </ul>	At least 400 g or 3 portions of vegetables (for calcium supply, select dark green vegetables frequently)
<b>Group 3:</b> Fruit (including nuts)	daily <ul style="list-style-type: none"> <li>At least 2 portions (250 g) fruit</li> </ul>	At least 300 g or 2 portions fruit daily, supplement fresh fruit with maximally 50 g dry fruit and fruit juices; Nuts and seeds: 30–60 g daily (in particular, almonds and sesame supply high levels of calcium)
<b>Group 4:</b> Milk and dairy products	daily <ul style="list-style-type: none"> <li>200–250 g low fat milk and dairy products</li> <li>2 slices (50–60 g) low fat cheese</li> </ul>	
<b>Group 5:</b> Meat, meat products, fish and eggs	weekly <ul style="list-style-type: none"> <li>300–600 g lean meat (prepared) and lean meat products</li> <li>1 portion (80–150 g) low fat saltwater fish (prepared) and</li> <li>1 portion (70 g) fatty saltwater fish (prepared) and up to 3 eggs (including processed eggs)</li> </ul>	not applicable; instead: legumes such as peas, beans, chickpeas and lentils: 1–2 meals per week and soya products (soya milk, soya yoghurt, tofu, tempeh, etc. ...) and other protein sources (e.g. seitan): 50–150 g/day
<b>Group 6:</b> Oils and fats	daily <ul style="list-style-type: none"> <li>10–15 g oil (e.g. rape, walnut or soya oil) and</li> <li>15–30 g margarine or butter</li> </ul>	unprocessed vegetable oils and fats: 2–4 tablespoons per day – best sources of n-3 fatty acids are rape, flaxseed and walnut oil
<b>Group 7:</b> Drinks	daily <ul style="list-style-type: none"> <li>about 1.5 L, preferably low energy or energy free drinks</li> </ul>	calcium-rich water and other alcohol free, low calorie drinks: daily 1–2 L
<b>Additional comments</b>	<ul style="list-style-type: none"> <li>Iodinated and fluoridated table salt is to be used sparingly.</li> <li>A unique characteristic of vitamin D is that it is not only obtained from food, but that the human body can also produce vitamin D itself if the skin is exposed to sunlight (UVB).</li> </ul>	<ul style="list-style-type: none"> <li>Sunlight (at least 15 min./day) for vitamin D</li> <li>In a vegan diet, the reliable supply of vitamin B<sub>12</sub> and D should be ensured by appropriate dietary supplements.</li> </ul>

Tab. 3: Food groups in the wholesome nutrition (mixed diet) and in the vegan nutrition

rian food pyramid emphasises dark green leafy vegetables and dried fruits. In addition, **cereals** in the form of whole-grain products and **potatoes** provide an additional basis and should be a component of meals. The recommendations for **oils, fats** and **drinks** are almost identical (butter and spreadable fat containing butter are excluded from a vegan diet). Although the food groups milk and dairy products and meat and meat products, as well as fish and eggs, are omitted from a vegan diet, the food-related recommendations for a vegan diet in the Giessen vegetarian food pyramid lay more emphasis on

**legumes, nuts and oil seeds**, as sources of protein, B vitamins, zinc and iron, as well as the use of fortified foods (e.g. soya milk with calcium). If fortified foods are not consumed or are unavailable, intake of various dietary supplements is recommended (vitamin B<sub>12</sub>, vitamin D, calcium). Measures to improve nutrient content (e.g. germination of cereals) and the bioavailability of nutrients from plant-based foods are additional strategies that should be recommended to improve nutrient supply in vegan nutrition. In international publications, it is often stated that **fortified foods**

may supply critical nutrients in vegan nutrition. This statement is essentially based on North America, where more fortified foods are on the market than are in Germany. A current survey on fortified foods available in different branches of food retailers in Germany (Bonn) permits the conclusion that in Germany fortified foods may make a noteworthy contribution to nutrient intake too; e.g. the reference value for riboflavin intake can be reached with 200 mL of a multivitamin juice. On the other hand, the added form and the bioavailability of the nutrients from these foods are

often unclear and the added quantities may be very variable within a single product group [72].

### International and national statements and recommendations

The Academy of Nutrition and Dietetics<sup>4</sup> [24]<sup>5</sup> takes the position that an appropriately planned vegan diet that includes dietary supplements and fortified foods is nutritionally adequate and is appropriate for individuals during all stages of the lifecycle, including pregnant and lactating women. This position is supported by scientific societies in other countries, including the National Health and Medical Research Council in the nutrition recommendations for Australia [69], the Portuguese National Programme for the Promotion of a Healthy Diet [74] and – for adults – the British Nutrition Foundation [25].

The Canadian Paediatric Society [75] also states that a well-planned vegan diet, including dietary supplements, can cover the nutrient requirements in children and adolescents, if adequate energy intake is ensured. **In the opinion of the British Nutrition Foundation [25] a well-planned, balanced vegetarian or vegan diet can be nutritionally adequate. More extreme diets, such as strict macrobiotic and raw food diets, are often low in energy and a range of micronutrients, making them wholly inadequate and inappropriate for children.** Moreover, the Portuguese National Programme for the Promotion of a Healthy Diet [74] recommends that breastfeeding for infants on a vegan diet should be extended beyond the recommended period of six months until 2 years of age during the food diversification process. In this way, it could be ensured that infants and toddlers received adequate supplies of high

quality milk protein.

The network „Healthy Start – Young Family Network“ states that a vegan diet is unsuitable for infants and toddlers, as it does not ensure their nutrient supplies. The network emphasises the risks for the child’s development, as well as the necessity of medical advice and taking dietary supplements [76–79]. Moreover, the recommendations of the Nutrition Committee of the German Society of Paediatrics and Adolescent Medicine rejects a vegan diet for healthy infants, unless dietary supplements are taken [68].

---

<sup>4</sup>until 2012 known as the American Dietetic Association (ADA) [24]

<sup>5</sup>The 2015 Position of the Academy of Nutrition and Dietetics: Vegetarian Diets [73] was withdrawn for revision during the preparation of this position paper. We therefore considered the 2009 Position [24].

## Conclusion: The DGE's position

An adequate intake of nutrients (see reference values for nutrient intake [30]) can be assured by a varied and diverse selection of foods. Any diet that does not lead to the intake of adequate levels of essential nutrients and energy is unfavourable.

The DGE recommends a diet that includes all groups of foods in the nutrition circle – including animal products. In other words, the DGE recommends a wholesome diet in the form of a mixed diet that largely consists of plant-based foods and, to a lesser extent, of animal foods, including fish, meat and meat products [80, 81].

The DGE also considers that pesco and ovo-lacto vegetarian diets are suitable for healthy persons in the long term, if appropriate alternatives are considered to optimize the nutrient intake. Special care is needed for groups with special requirements for nutrient supply, e.g. pregnant women, lactating women, infants and toddlers.

On a vegan diet, it is difficult or impossible to ensure adequate supply of some nutrients. The most critical nutrient is vitamin B<sub>12</sub>. Other potentially critical nutrients on a vegan diet include protein resp. indispensable amino acids and long-chain n-3 fatty acids (EPA and DHA), other vitamins (riboflavin, vitamin D) and minerals (calcium, iron, iodine, zinc and selenium). A vegan diet is only able to fulfil requirements if the following points are considered. Persons on a vegan diet should

- take a vitamin B<sub>12</sub> supplement in the long term and have their supplies of vitamin B<sub>12</sub> regularly checked by a physician;
- select very specifically nutrient

dense foods and fortified foods, in order to ensure supply of nutrients, particularly critical nutrients (♦ Table 2);

- possibly have the supplies of other critical nutrients regularly checked by a physician; if there is a definite or possible nutritional deficiency, the nutrition should be adjusted and the critical nutrients should be added – either in dietary supplements or fortified foods, until the nutrient deficiency has been corrected;
- consult a qualified nutrition counsellor [82] for receiving advice.

The risk of nutrient undersupply or a nutritional deficiency is greater in persons in sensitive phases of life, such as pregnancy, lactation and in infants, children and adolescents taking or being given a vegan diet, than in healthy adults on a vegan diet. With some nutrients, a vegan diet without fortified foods or dietary supplements leads to inadequate intake, which may have considerable unfavourable consequences for health. Since rejecting any animal foods increases the risk of nutrient deficiencies and thus of health disorders, a vegan diet is not recommended by the DGE during pregnancy or lactation, or for children or adolescents of any age. It is essential that persons who nevertheless decide to adhere to a vegan diet should note the above points.

### Important comment:

This statement has been carefully checked for its content by the publisher; however, a guarantee for the content cannot be accepted. Neither the publisher, nor the editors are liable for any personal injury or damage to property.

### Acknowledgement

The authors wish to thank Prof. Ulrike ARENS-AZEVEDO, Prof. Dr. Helmut ERBERSDOBLER, Prof. Dr. Hans HAUNER, Prof. Dr. Michael KRAWINKEL, Prof. Dr. Monika NEUHÄUSER-BERTHOLD, Prof. Dr. Hildegard PRZYREMBEL, Prof. Dr. Gerhard RECHKEMMER, Prof. Dr. Gabriele STANGL, Prof. Dr. Peter STEHLE, Prof. Dr. Karl-Heinz WAGNER, Prof. Dr. Gertrud WINKLER and Prof. Dr. Günther WOLFRAM for critically reviewing the manuscript and for valuable discussions.

### Corresponding Author:

Dr. Margrit Richter  
E-Mail: richter@dge.de

Dr. Margrit Richter<sup>1</sup>  
Prof. Dr. Heiner Boeing<sup>2</sup>  
Dr. Dorle Grünewald-Funk<sup>3</sup>  
Prof. Dr. Helmut Heseker<sup>4</sup>  
Prof. Dr. Anja Kroke<sup>5</sup>  
Dr. Eva Leschik-Bonnet<sup>1</sup>  
Dr. Helmut Oberritter<sup>1</sup>  
Dr. Daniela Strohm<sup>1</sup>  
Prof. Dr. Bernhard Watzl<sup>6</sup>

<sup>1</sup> Deutsche Gesellschaft für Ernährung e. V. (DGE), Godesberger Allee 18, 53175 Bonn

<sup>2</sup> Deutsches Institut für Ernährungsforschung (DIfE), Arthur-Scheunert-Allee 114–116, 14558 Nuthetal

<sup>3</sup> kompetenz für kommunikation public health ernährung, PF 02 24 50, 10126 Berlin

<sup>4</sup> Institut für Ernährung, Konsum und Gesundheit, Universität Paderborn, Warburger Str. 100, 33098 Paderborn

<sup>5</sup> Hochschule Fulda, Fachbereich Oecotrophologie, Leipziger Str. 123, 36037 Fulda

<sup>6</sup> Max Rubner-Institut, Bundesforschungsinstitut für Ernährung und Lebensmittel, Institut für Physiologie und Biochemie der Ernährung, Haid-und-Neu-Str. 9, 76131 Karlsruhe

### Conflict of Interest

The authors declare no conflict of interest.



References

1. Kerschke-Risch P (2015) Vegan diet: motives, approach and duration. Initial results of a quantitative sociological study. *Ernährungs Umschau* 62: 98–103
2. Leitzmann C, Keller M. *Vegetarische Ernährung*. 3., aktual. Aufl., Ulmer, Stuttgart (2013)
3. Piccoli GB, Clari R, Vigotti FN et al. (2015) Vegan-vegetarian diets in pregnancy: danger or panacea? A systematic narrative review. *BJOG* 122: 623–633
4. Deutsche Gesellschaft für Ernährung (DGE) (2013) Flexitarier – die flexiblen Vegetarier. *DGEInfo* 146–149
5. Bouvard V, Loomis D, Guyton KZ et al. (2015) Carcinogenicity of consumption of red and processed meat. *Lancet Oncol* 16: 1599–1600
6. Boeing H. Prävention durch Ernährung. In: Deutsche Gesellschaft für Ernährung (Hg). 12. Ernährungsbericht 2012, Bonn (2012), S. 319–354
7. World Cancer Research Fund (WCRF), American Institute for Cancer Research (AICR) (Hg). *Food, nutrition, physical activity, and the prevention of cancer: a global perspective*. Washington DC (2007)
8. Boeing H, Bechthold A, Bub A et al. (2012) Critical review: vegetables and fruit in the prevention of chronic diseases. *Eur J Clin Nutr* 51: 637–663
9. Deutsche Gesellschaft für Ernährung (DGE) (Hg). *Kohlenhydratzufuhr und Prävention ausgewählter ernährungsmitbedingter Krankheiten – Evidenzbasierte Leitlinie*. Bonn (2011) URL: [www.dge.de/wissenschaft/leitlinien/leitlinie-kohlenhydrate/](http://www.dge.de/wissenschaft/leitlinien/leitlinie-kohlenhydrate/) Zugriff 11.02.16
10. Dinu M, Abbate R, Gensini GF et al. (2016) Vegetarian, vegan diets and multiple health outcomes: a systematic review with meta-analysis of observational studies. *Crit Rev Food Sci Nutr* [Epub ahead of print]
11. Kwok CS, Umar S, Myint PK et al. (2014) Vegetarian diet, Seventh Day Adventists and risk of cardiovascular mortality: A systematic review and meta-analysis. *Int J Cardiol* 176: 680–686
12. Appleby PN, Crowe FL, Bradbury KE et al. (2016) Mortality in vegetarians and comparable nonvegetarians in the United Kingdom. *Am J Clin Nutr* 103: 218–230
13. Clarys P, Deliens T, Huybrechts I et al. (2014) Comparison of nutritional quality of the vegan, vegetarian, semi-vegetarian, pesco-vegetarian and omnivorous diet. *Nutrients* 6: 1318–1332
14. Kristensen NB, Madsen ML, Hansen TH et al. (2015) Intake of macro- and micronutrients in Danish vegans. *Nutr J* 14: 115
15. Orlich MJ, Jaceldo-Siegl K, Sabaté J et al. (2014) Patterns of food consumption among vegetarians and non-vegetarians. *Br J Nutr* 112: 1644–1653
16. Barr SI, Broughton TM (2000) Relative weight, weight loss efforts and nutrient intakes among health-conscious vegetarian, past vegetarian and nonvegetarian women ages 18 to 50. *J Am Coll Nutr* 19: 781–788
17. Davey GK, Spencer EA, Appleby PN et al. (2003) EPIC-Oxford: lifestyle characteristics and nutrient intakes in a cohort of 33 883 meat-eaters and 31 546 non meat-eaters in the UK. *Public Health Nutr* 6: 259–269
18. Draper A, Lewis J, Malhotra N et al. (1993) The energy and nutrient intakes of different types of vegetarian: a case for supplements? *Br J Nutr* 69: 3–19
19. Deriemaeker P, Alewaeters K, Hebbelinck M et al. (2010) Nutritional status of Flemish vegetarians compared with non-vegetarians: a matched samples study. *Nutrients* 2: 770–780
20. Gilsing, AMJ, Crowe FL, Lloyd-Wright Z et al. (2010) Serum concentrations of vitamin B<sub>12</sub> and folate in British male omnivores, vegetarians and vegans: results from a cross-sectional analysis of the EPIC-Oxford cohort study. *Eur J Clin Nutr* 64: 933–939
21. Krajcovicová-Kudláčková M, Simoncic R, Babinská K et al. (1995) Selected vitamins and trace elements in blood of vegetarians. *Ann Nutr Metab* 39: 334–339
22. Majchrzak D, Singer I, Männer M et al. (2006) B-vitamin status and concentrations of homocysteine in Austrian omnivores, vegetarians and vegans. *Ann Nutr Metab* 50: 485–491
23. Rizzo NS, Jaceldo-Siegl K, Sabate J et al. (2013) Nutrient profiles of vegetarian and nonvegetarian dietary patterns. *J Acad Nutr Diet* 113: 1610–1619
24. Craig WJ, Reed Mangels A, American Dietetic Association (2009) Position of the American Dietetic Association: vegetarian diets. *J Am Diet Assoc* 109: 1266–1282
25. Phillips F (2005) Vegetarian nutrition. *Nutrition Bulletin* 30: 132–167
26. Jones G. Vitamin D. In: Ross AC, Caballero B, Cousins RJ et al. (Hg). *Modern nutrition in health and disease*. 11. Aufl., Lippincott Williams & Wilkins, Philadelphia (2014), S. 278–292
27. Zittermann A, Gummert JF (2010) Non-classical vitamin D actions. *Nutrients* 2: 408–425
28. Krems C, Walter C, Heuer T et al. *Lebensmittelverzehr und Nährstoffzufuhr – Ergebnisse der Nationalen Verzehrsstudie II*. In: Deutsche Gesellschaft für Ernährung (Hg). 12. Ernährungsbericht 2012. Bonn (2012), S. 40–85
29. Craig WJ (2009) Health effects of vegan diets. *Am J Clin Nutr* 89: 1627S–1633S
30. Deutsche Gesellschaft für Ernährung (DGE), Österreichische Gesellschaft für Ernährung (ÖGE), Schweizerische Gesellschaft für Ernährung (SGE) (Hg). *Referenzwerte für die Nährstoffzufuhr*. 2. Aufl., 1. Ausg., Bonn (2015)
31. Maqdry E, Lisowska A, Grebowiec P et al. (2013) The impact of vegan diet on B-12 status in healthy omnivores: five-year prospective study. *Acta Sci Pol Technol Aliment* 11: 209–212
32. Pawlak R, Parrott SJ, Raj S et al. (2013) How prevalent is vitamin B(12) deficiency among vegetarians? *Nutr Rev* 71: 110–117
33. Pawlak R, Lester SE, Babatunde T (2014) The prevalence of cobalamin deficiency among vegetarians assessed by serum vitamin B<sub>12</sub>: a review of literature. *Eur J Clin Nutr* 68: 541–548
34. Waldmann A, Koschizke JW, Leitzmann C et al. (2004) Homocysteine and cobalamin status in German vegans. *Public Health Nutr* 7: 467–472
35. Hahn A, Waldmann A (2004) *Gesund mit reiner Pflanzenkost?* Unimagazin Hannover: 6–9
36. Obersby D, Chappell DC, Dunnett A et al. (2013) Plasma total homocysteine status of vegetarians compared with omnivores: a systematic review and meta-analysis. *Br J Nutr* 109: 785–794
37. Carmel R. Cobalamin (Vitamin B<sub>12</sub>). In: Ross AC, Caballero B, Cousins RJ et al. (Hg). *Modern nutrition in health and disease*. 11. Aufl., Lippincott Williams & Wilkins, Philadelphia (2014), S. 369–389
38. Black MM (2008) Effects of vitamin B<sub>12</sub> and folate deficiency on brain development in children. *Food Nutr Bull* 29: S126–S131

39. Finkelstein JL, Layden AJ, Stover PJ (2015) Vitamin B-12 and perinatal health. *Adv Nutr* 6: 552–563
40. Molloy AM, Kirke PN, Brody LC et al. (2008) Effects of folate and vitamin B<sub>12</sub> deficiencies during pregnancy on fetal, infant, and child development. *Food Nutr Bull* 29: S101–S111; discussion S112–S115
41. Molloy AM, Kirke PN, Troendle JF et al. (2009) Maternal vitamin B<sub>12</sub> status and risk of neural tube defects in a population with high neural tube defect prevalence and no folic acid fortification. *Pediatrics* 123: 917–923
42. Antony AC (2003) Vegetarianism and vitamin B-12 (cobalamin) deficiency. *Am J Clin Nutr* 78: 3–6
43. Casella EB, Valente M, de Navarro JM et al. (2005) Vitamin B<sub>12</sub> deficiency in infancy as a cause of developmental regression. *Brain Dev* 27: 592–594
44. Ciani F, Poggi GM, Pasquini E et al. (2000) Prolonged exclusive breast-feeding from vegan mother causing an acute onset of isolated methylmalonic aciduria due to a mild mutase deficiency. *Clin Nutr* 19: 137–139
45. Dror DK, Allen LH (2008) Effect of vitamin B<sub>12</sub> deficiency on neurodevelopment in infants: current knowledge and possible mechanisms. *Nutr Rev* 66: 250–255
46. Lücke T, Korenke GC, Poggenburg I et al. (2007) Mütterlicher Vitamin-B<sub>12</sub>-Mangel: Ursache neurologischer Symptomatik im Säuglingsalter. *Z Geburtshilfe Neonatol* 211: 157–161
47. Schlapbach LJ, Schütz B, Nuoffer JM et al. (2007) Floppy Baby mit makrozytärer Anämie und veganischer Mutter. *Praxis* 96: 1309–1314
48. Shinwell ED, Gorodischer R (1982) Totally vegetarian diets and infant nutrition. *Pediatrics* 70: 582–586
49. Sklar R (1986) Nutritional vitamin B<sub>12</sub> deficiency in a breast-fed infant of a vegan-diet mother. *Clin Pediatr (Phila)* 25: 219–221
50. Specker BL, Black A, Allen L et al. (1990) Vitamin B-12: low milk concentrations are related to low serum concentrations in vegetarian women and to methylmalonic aciduria in their infants. *Am J Clin Nutr* 52: 1073–1076
51. Stabler SP, Allen RH (2004) Vitamin B<sub>12</sub> deficiency as a worldwide problem. *Annu Rev Nutr* 24: 299–326
52. Stötter M, Mayrhofer H (1996) Veganische Ernährung: Neurologische Symptomatik, schwere Entwicklungs- und Gedeihstörung bei Säuglingen und Kleinkindern durch Vitamin B<sub>12</sub>-Mangel. *Aktuel Ernähr Med* 21: 4–7
53. Wagnon J, Cagnard B, Bridoux-Henno L et al. (2005) Allaitement maternel et végétalisme. *J Gynecol Obstet Biol Reprod (Paris)* 34: 610–612
54. Weiss R, Fogelman Y, Bennett M (2004) Severe vitamin B<sub>12</sub> deficiency in an infant associated with a maternal deficiency and a strict vegetarian diet. *J Pediatr Hematol Oncol* 26: 270–271
55. Green R (2011) Indicators for assessing folate and vitamin B-12 status and for monitoring the efficacy of intervention strategies. *Am J Clin Nutr* 94 (Suppl): 666S–672S
56. Shaikh MG, Anderson JM, Hall SK et al. (2003) Transient neonatal hypothyroidism due to a maternal vegan diet. *J Pediatr Endocrinol Metab* 16: 111–113
57. Deutsche Gesellschaft für Ernährung (DGE) (Hg). Fettzufuhr und Prävention ausgewählter ernährungsmittbedingter Krankheiten – Evidenzbasierte Leitlinie. Bonn (2015) URL: [www.dge.de/index.php?id=207#c313](http://www.dge.de/index.php?id=207#c313) Zugriff 11.02.16
58. Strohm D, Boeing H, Leschik-Bonnet E et al. for the German Nutrition Society (2016) Salt intake in Germany, health consequences, and resulting recommendations for action. A scientific statement from the German Nutrition Society (DGE). *Ernährungs Umschau* 63: 62–70
59. Radnitz C, Beezhold B, DiMatteo J (2015) Investigation of lifestyle choices of individuals following a vegan diet for health and ethical reasons. *Appetite* 90: 31–36
60. Hauner H (2015) Der ultimative Hype. *MMW Fortschritte der Medizin* 157: 40
61. Leitzmann C (2013) Fleischersatz – rein pflanzlich. *UIG-Forum* 30: 296–299
62. Verbraucherzentrale Hamburg (Hg). Nix mit Tieren, aber gesund? 2015 URL: [www.vzhh.de/ernaehrung/334670/nix-mit-tieren-aber-gesund.aspx](http://www.vzhh.de/ernaehrung/334670/nix-mit-tieren-aber-gesund.aspx) Zugriff 11.02.16
63. Öko-Test (Hg). Vegane/Vegetarische Lebensmittel. So ja! 2014. URL: [www.oekotest.de/cgi/index.cgi?artnr=104240&bernr=04&gartnr=1&suche=fleischersatz](http://www.oekotest.de/cgi/index.cgi?artnr=104240&bernr=04&gartnr=1&suche=fleischersatz) Zugriff 25.06.15
64. Vandenplas Y, Castrellon PG, Rivas R et al. (2014) Safety of soya-based infant formulas in children. *Br J Nutr* 111: 1340–1360
65. Bundesinstitut für Risikobewertung (BfR). Säuglingsnahrung aus Sojaweiß ist kein Ersatz für Kuhmilchprodukte. (2007) URL: [www.bfr.bund.de/cm/343/saeuglingsnahrung\\_aus\\_sojaeiweiss\\_ist\\_kein\\_ersatz\\_fuer\\_kuhmilchprodukte.pdf](http://www.bfr.bund.de/cm/343/saeuglingsnahrung_aus_sojaeiweiss_ist_kein_ersatz_fuer_kuhmilchprodukte.pdf) Zugriff 11.02.16
66. Koletzko B, Brönstrup A, Cremer M et al. (2013) Säuglingsernährung und Ernährung der stillenden Mutter. *Monatsschr Kinderheilkd* 161: 237–246
67. Koletzko B (2006) Stellungnahme zur Verwendung von Säuglingsnahrungen auf Sojaweißbasis. *Monatsschr Kinderheilkd* 154: 913–916
68. Bühner C, Genzel-Boroviczeny O, Jochum F et al. (2014) Ernährung gesunder Säuglinge. *Monatsschr Kinderheilkd* 162: 527–538
69. National Health and Medical Research Council (Hg). Eat for health. Australian dietary guidelines. National Health and Medical Research Council, Canberra (2013)
70. United States Department of Agriculture (USDA). Scientific Report of the 2015 Dietary Guidelines Advisory Committee. (2015) URL: [www.health.gov/dietaryguidelines/2015-scientific-report/](http://www.health.gov/dietaryguidelines/2015-scientific-report/) Zugriff 11.02.16
71. United States Department of Agriculture (USDA) (Hg). Healthy eating for vegetarians. 10 tips for vegetarians. (2011) URL: [www.choosemyplate.gov/sites/default/files/tentips/DGTipsheet8HealthyEatingForVegetarians.pdf](http://www.choosemyplate.gov/sites/default/files/tentips/DGTipsheet8HealthyEatingForVegetarians.pdf) Zugriff 03.02.16
72. Marczykowski FJ. Ist eine Deckung der D-A-CH-Referenzwerte für kritische Nährstoffe bei einer veganen Ernährung über natürliche und angereicherte Lebensmittel möglich? Bachelorarbeit. Bonn (2015)
73. Cullum-Dugan D, Pawlak R (2015) Position of the academy of nutrition and dietetics: vegetarian diets. *J Acad Nutr Diet* 115: 801–810
74. National Programme for the Promotion of a Healthy Diet, Direção-Geral da Saúde (Hg). Guidelines for a healthy vegetarian diet. Lissabon (2015)
75. Amit M (2010) Vegetarian diets in children and adolescents. *Paediatr Child Health* 15: 303–314 [reaffirmed 2014]
76. Koletzko B, Armbruster M, Bauer C et al. (2013) Ernährung und Bewegung im Kleinkindalter. *Monatsschr Kinderheilkd* 161: 1187–1200
77. Koletzko B, Bauer CP, Bung P et al. (2012)

- Ernährung in der Schwangerschaft – Teil 1. *Dtsch med Wochenschr* 137: 1309–1313
78. Koletzko B, Bauer C, Bung P et al. (2012) Ernährung in der Schwangerschaft – Teil 2. Handlungsempfehlungen des Netzwerks „Gesund ins Leben – Netzwerk Junge Familie“. *Dtsch med Wochenschr* 137: 1366–1372
79. Koletzko B, Brönstrup A, Cremer M et al. (2010) Säuglingsernährung und Ernährung der stillenden Mutter. Handlungsempfehlungen – Ein Konsensuspapier im Auftrag des bundesweiten Netzwerk Junge Familie. *Monatschr Kinderheilkd* 158: 679–689
80. Deutsche Gesellschaft für Ernährung (DGE) (Hg). *Vollwertig essen und trinken nach den 10 Regeln der DGE*. 25., überarb. Aufl., Bonn (2013)
81. Deutsche Gesellschaft für Ernährung (DGE). *Lebensmittelbezogene Ernährungsempfehlungen in Deutschland*. DGE, Bonn (2014)
82. Koordinierungskreis zur Qualitätssicherung in der Ernährungsberatung und Ernährungsbildung (Hg). *Rahmenvereinbarung zur Qualitätssicherung in der Ernährungsberatung und Ernährungsbildung in Deutschland*. In der Fassung vom 16.06.2014 (2015) URL: [www.dge.de/fileadmin/public/doc/fb/14-06-16-KoKreis-EB-RV.pdf](http://www.dge.de/fileadmin/public/doc/fb/14-06-16-KoKreis-EB-RV.pdf) Zugriff 15.02.16
83. Young VR, Pellett PL (1994) Plant proteins in relation to human protein and amino acid nutrition. *Am J Clin Nutr* 59: 1203S–1212S
84. World Health Organization (WHO). *Protein and amino acid requirements in human nutrition*. WHO, Genf (2007)
85. Marsh KA, Munn EA, Baines SK (2013) Protein and vegetarian diets. *Med J Aust* 199(Suppl): S7–S10
86. Sanders TA (1999) The nutritional adequacy of plant-based diets. *Proc Nutr Soc* 58: 265–269
87. Krajcovicova-Kudlackova M, Simoncic R, Bederova A et al. (1997) Influence of vegetarian and mixed nutrition on selected haematological and biochemical parameters in children. *Nahrung* 41: 311–314
88. Boye J, Wijesinha-Bettoni R, Burlingame B (2012) Protein quality evaluation twenty years after the introduction of the protein digestibility corrected amino acid score method. *Br J Nutr* 108(Suppl 2): S183–S211
89. Tome D (2012) Criteria and markers for protein quality assessment – a review. *Br J Nutr* 108(Suppl 2): S222–S229
90. Matthews DE. *Proteins and amino acids*. In: Ross AC, Caballero B, Cousins RJ et al. (Hg). *Modern nutrition in health and disease*. 11. Aufl., Lippincott Williams & Wilkins, Philadelphia (2014), S. 3–35
91. Brenna JT (2002) Efficiency of conversion of alpha-linolenic acid to long chain n-3 fatty acids in man. *Curr Opin Clin Nutr Metab Care* 5: 127–132
92. Geppert J, Kraft V, Demmelmair H et al. (2006) Microalgal docosahexaenoic acid decreases plasma triacylglycerol in normolipidaemic vegetarians: a randomised trial. *Br J Nutr* 95: 779–786
93. Lane K, Derbyshire E, Li W et al. (2014) Bioavailability and potential uses of vegetarian sources of omega-3 fatty acids: a review of the literature. *Crit Rev Food Sci Nutr* 54: 572–579
94. Sarter B, Kelsey KS, Schwartz TA et al. (2015) Blood docosahexaenoic acid and eicosapentaenoic acid in vegans: Associations with age and gender and effects of an algal-derived omega-3 fatty acid supplement. *Clin Nutr* 34: 212–218
95. Durchführungsbeschluss der Kommission vom 14. Juli 2014 zur Genehmigung des Inverkehrbringens von Öl aus der Mikroalge *Schizochytrium sp.* als neuartige Lebensmittelzutat im Sinne der Verordnung (EG) Nr. 258/97 des Europäischen Parlaments und des Rates und zur Aufhebung der Entscheidungen 2003/427/EG und 2009/778/EG. *Amtsblatt der Europäischen Union* (2009) 55–58
96. Entscheidung der Kommission vom 21. Oktober 2009 über die Erweiterung der Anwendungen von Algenöl aus der Mikroalge *Ulkenia sp.* als neuartige Lebensmittelzutat im Sinne der Verordnung (EG) Nr. 258/97 des Europäischen Parlaments und des Rates. *Amtsblatt der Europäischen Union* (2009) 54–55
97. Watanabe F, Yabuta Y, Bito T et al. (2014) Vitamin B<sub>12</sub>-containing plant food sources for vegetarians. *Nutrients* 6: 1861–1873
98. Croft MT, Lawrence AD, Raux-Deery E et al. (2005) Algae acquire vitamin B<sub>12</sub> through a symbiotic relationship with bacteria. *Nature* 438: 90–93
99. Kittaka-Katsura H, Fujita T, Watanabe F et al. (2002) Purification and characterization of a corrinoid compound from *Chlorella* tablets as an algal health food. *J Agric Food Chem* 50: 4994–4997
100. Watanabe F, Takenaka S, Katsura H et al. (1999) Dried green and purple lavers (Nori) contain substantial amounts of biologically active vitamin B<sub>12</sub> but less of dietary iodine relative to other edible seaweeds. *J Agric Food Chem* 47: 2341–2343
101. Watanabe F (2007) Vitamin B<sub>12</sub> sources and bioavailability. *Exp Biol Med* (Maywood) 232: 1266–1274
102. Watanabe F, Yabuta Y, Tanioka Y et al. (2013) Biologically active vitamin B<sub>12</sub> compounds in foods for preventing deficiency among vegetarians and elderly subjects. *J Agric Food Chem* 61: 6769–6775
103. Hurrell R, Egli I (2010) Iron bioavailability and dietary reference values. *Am J Clin Nutr* 91: 1461S–1467S
104. Bundesinstitut für Risikobewertung (BfR). *Gesundheitliche Risiken durch zu hohen Jodgehalt in getrockneten Algen*. (2007) URL: [www.bfr.bund.de/cm/343/gesundheitliche\\_risiken\\_durch\\_zu\\_hohen\\_jodgehalt\\_in\\_getrockneten\\_algen.pdf](http://www.bfr.bund.de/cm/343/gesundheitliche_risiken_durch_zu_hohen_jodgehalt_in_getrockneten_algen.pdf) Zugriff 11.02.16

DOI: 10.4455/eu.2016.021